### Gö-VIP-04: Prof. Dr. Mikael Simons

Klinik für Neurologie

**Originalpublikation**: "Myelin Membrane Wrapping of CNS Axons by PI(3,4,5)P3-Dependent Polarized Growth at the Inner Tongue." In: Cell 156, 277–290, January 16, 2014, Autoren: Nicolas Snaidero (1,2) Wiebke Möbius, Tim Czopka, Liesbeth H.P. Hekking, Cliff Mathisen, Dick Verkleij, Sandra Goebbels, Julia Edgar, Doron Merkler, David A. Lyons, Klaus-Armin Nave, and Mikael Simons (1,2)\*

(1) Max Planck Institute of Experimental Medicine, Cellular Neuroscience, Hermann-Rein-Straße, 3, 37075 Göttingen, Germany

(2) Department of Neurology, University of Göttingen, Robert-Koch-Straße, 40, 37075 Göttingen, Germany

\*Corresponding Author



Abb.: Prof. Dr. Mikael Simons, Prof. Dr. Nikolas Snaidero (von links nach rechts)

# Zusammenfassung des wissenschaftlichen Inhalts (Prof. Dr. Mikael Simons)

Myelin ist eine multilamelläre Membran, die von Oligodendrozyten im ZNS zur schnellen Erregungsfortleitung generiert wird. Große Teile des Gehirns bestehen aus Myelin, insbesondere die tiefer liegende weiße Substanz des Gehirns, die aus Millionen langer, von weißlichem Myelin umhüllten Axonen besteht.

Mikael Simons und sein Team (MPI für Experimentelle Medizin und der Abteilung für Neurologie der Universität Göttingen) gingen der Frage nach wie Myelin gebildet wird. Fest steht, dass für die Bildung des Myelin Gliazellen, sogenannte Oligodendrozyten verantwortlich sind, deren Zellfortsätze sich mehrfach um die Axone winden.

Doch wie wickeln Oligodendrozyten ihre Membran um das Axon und wie wird das Myelin-Wachstum gesteuert? Um dies herauszufinden, haben die Wissenschaftler die Myelinbildung in seiner Entwicklung untersucht und jetzt die Wachstumszone identifizieren können. Dabei fanden sie heraus, dass sich die innerste Membranschicht um das Axon windet und sich gleichzeitig entlang des Axons nach außen bewegt. Durch hochauflösende und dreidimensionale elektronenmikroskopische Aufnahmen konnten die Wissenschafter feststellen, dass sich während der Entwicklung tunnelähnliche Transportwege in der Myelinmembran bilden. Entlang dieser Transportwege wird das neu synthetisierte Material befördert, bevor es schließlich in der sich ausbreitenden Membran eingebaut wird.

Wenn der Wachstumsprozess abgeschlossen ist, schließen sich die Tunnel im Myelin wieder und die Materialzufuhr kommt zum Erliegen. Dies geschieht bei Menschen erst sehr spät. Neugeborene enthalten nur wenige Bereiche mit umhüllten Fasern. In Kindheit und Jugend nimmt das Ausmaß der Myelinisierung kontinuierlich zu. Erst im Jungendalter sind alle Hirnregionen voll versorgt. Bisher ist man davon ausgegangen, dass nach Abschluss der Myelinbildung die Membran nicht weiterwachsen kann. Die Wissenschaftler konnten zeigen, dass durch die künstliche Erhöhung eines wichtigen Signalsstoffes (Phosphatidylinositol (3,4,5)-triphosphate), das Wachstum wieder in Gang gesetzt werden kann, da sich dadurch die Tunnel zum Transport der Membran im Myelin neu öffnen. Damit das Netzwerk der Nerven intakt bleibt, muss also immer wieder neues Myelin nachgeliefert werden. Diese Mechanismen sind nicht nur zum Verständnis der normalen Gehirnentwicklung von Bedeutung, sondern könnten auch bei der Entwicklung von neuen therapeutischen Strategien der Myelin-Reparatur bei Erkrankungen wie der Multiplen Sklerose hilfreich sein.

### WEITERE INFORMATIONEN:

**Prof. Dr. Mikael Simons** Einrichtung: <u>Klinik für Neurologie</u> Telefon: 0551-39-6715 / -22685 E-Mail: <u>msimons@gwdg.de</u>

Weitere Links

- PubMed
- More information and download the publication



Cell

# Myelin Membrane Wrapping of CNS Axons by PI(3,4,5)P3-Dependent Polarized Growth at the Inner Tongue

Nicolas Snaidero,<sup>1,2</sup> Wiebke Möbius,<sup>3,12</sup> Tim Czopka,<sup>4,5,6,7</sup> Liesbeth H.P. Hekking,<sup>8</sup> Cliff Mathisen,<sup>8</sup> Dick Verkleij,<sup>8</sup> Sandra Goebbels,<sup>3</sup> Julia Edgar,<sup>3</sup> Doron Merkler,<sup>9,10,11</sup> David A. Lyons,<sup>4,5,6</sup> Klaus-Armin Nave,<sup>3</sup> and Mikael Simons<sup>1,2,\*</sup> <sup>1</sup>Max Planck Institute of Experimental Medicine, Cellular Neuroscience, Hermann-Rein-Strasse, 3, 37075 Göttingen, Germany

<sup>2</sup>Department of Neurology, University of Göttingen, Robert-Koch-Strasse, 40, 37075 Göttingen, Germany

<sup>3</sup>Department of Neurogenetics, Max Planck Institute of Experimental Medicine, Hermann-Rein-Strasse, 3, 37075 Göttingen, Germany <sup>4</sup>Centre for Neuroregeneration

<sup>5</sup>MS Society Centre for Translational Research

<sup>6</sup>Euan Mac Donald Centre for Motor Neurone Disease Research

<sup>7</sup>MRC Centre for Regenerative Medicine, University of Edinburgh, Edinburgh EH16 4UU, UK

<sup>8</sup>FEI Company, Achtseweg Noord 5, 5651 GG Eindhoven, The Netherlands

<sup>9</sup>Department of Pathology and Immunology, University of Geneva, 1211 Geneva, Switzerland

<sup>10</sup>Division of Clinical Pathology, Geneva University Hospital, 1211 Geneva, Switzerland

<sup>11</sup>Department of Neuropathology, University of Göttingen, 37075 Göttingen, Germany

<sup>12</sup>Center for Nanoscale Microscopy and Molecular Physiology of the Brain (CNMPB), 37075 Göttingen, Germany

\*Correspondence: msimons@gwdg.de

http://dx.doi.org/10.1016/j.cell.2013.11.044

### SUMMARY

Central nervous system myelin is a multilayered membrane sheath generated by oligodendrocytes for rapid impulse propagation. However, the underlying mechanisms of myelin wrapping have remained unclear. Using an integrative approach of live imaging, electron microscopy, and genetics, we show that new myelin membranes are incorporated adjacent to the axon at the innermost tongue. Simultaneously, newly formed layers extend laterally, ultimately leading to the formation of a set of closely apposed paranodal loops. An elaborated system of cytoplasmic channels within the growing myelin sheath enables membrane trafficking to the leading edge. Most of these channels close with ongoing development but can be reopened in adults by experimentally raising phosphatidylinositol-(3,4,5)triphosphate levels, which reinitiates myelin growth. Our model can explain assembly of myelin as a multilayered structure, abnormal myelin outfoldings in neurological disease, and plasticity of myelin biogenesis observed in adult life.

### INTRODUCTION

In the vertebrate nervous system, axons are ensheathed with myelin, which is one of the most remarkable and complex transformations of a plasma membrane (Jessen and Mirsky, 2005; Sherman and Brophy, 2005; Salzer et al., 2008). More than 60

years after the seminal discovery demonstrating that myelin is made by axon-associated glial cells, and not by the axon itself (Ben Geren, 1954), the molecular mechanisms by which the myelin sheath is wrapped around the axon are still largely unknown. This is due in part to the physical limitations of visualizing membrane dynamics at the nanometer scale and the time span involved (i.e., days in vivo). Even if it represents "textbook knowledge" that oligodendrocytes wrap myelin around an axon by steering a leading process that stays in close contact with the axon, we have almost no experimental data to substantiate this claim. Does the leading edge resemble a glial growth cone-like extension related to the one that drives axonal outgrowth in developing neurons? It has also become apparent that myelin is a dynamically active structure (Young et al., 2013) that can provide metabolic support to associated axons (Fünfschilling et al., 2012; Lee et al., 2012). However, it remains completely unclear how molecules reach the innermost myelin layer, i.e., passing through a multilamellar stack of membranes.

A number of different models have been proposed to explain how a myelin sheath might form in development. According to the "carpet crawler" model (Bunge et al., 1961, 1989), the oligodendrocyte forms a process that broadens and extends along the entire axonal segment (the future internode) before it makes one turn and moves underneath the growing sheet. However, at least in the CNS, several morphological features of myelin are incompatible with this model. In particular, it is clear from electron microscopic analysis that the number of myelin layers can vary along the length of a single myelin sheath during its formation (Knobler et al., 1976). Moreover, the molecular forces necessary to continuously displace myelin by newly made layers of membrane from underneath might be too high. Some of these shortcomings were reconciled in the "liquid croissant" (Sobottka



University of Edinburgh, Edinburgh EH16 4SB, UK

et al., 2011) and the "yo-yo" model (D. Colman, personal communication). In the latter case, myelin growth begins with a single glial process that, after making axonal contact, spirally encircles the future internode, followed by the lateral growth of the individual membrane layers. However, because of the physical limitations of imaging such processes, it has not been possible to experimentally support any of these models of myelin wrapping.

Here, by combining an array of techniques (in vivo live-imaging, serial block-face imaging by focused ion beam, high-pressure freezing, and a virus as a reporter), we show that myelin growth occurs by the consecutive wrapping of the inner tongue (leading edge) around the axon and the coordinated lateral extension of the individual layers of myelin. Cytoplasmic subdomains at the end of each myelin layer remain in close contact with the underlying axon moving laterally and around the axon toward the future node, where they eventually form a set of closely apposed "paranodal loops." Together, our data elucidate mechanisms of myelin sheath morphogenesis and its regulation throughout life.

### RESULTS

### Visualization of Nascent Myelin Sheath Morphogenesis In Vivo

In order to visualize the dynamics of myelin sheath formation in vivo, we included the zebrafish as a model organism, given its amenability for high-resolution in vivo imaging (Kirby et al., 2006). In order to resolve the initial wrapping events of CNS myelination, we utilized the Tg(nkx2.2a:meGFP) line, which expresses a membrane-tethered GFP in early myelinating oligodendrocytes (Kirby et al., 2006; Czopka et al., 2013). We imaged these early myelination events and plotted the fluorescence intensity profiles along the length of 32 short (less than 10  $\mu$ m in length) nascent myelin sheaths and found that fluorescence intensity was consistently higher in the central regions of the sheaths relative to their ends (Figures 1A-1C; Figure S1 available online). By electron microscopy (EM), we found that most of the myelin sheaths with less than four wraps were still in an uncompacted state (Figure S1). We hypothesized that the change in fluorescence intensity might reflect differences in the number of wraps of myelin membrane around the axon at different points along the myelin sheath. In order to address this possibility, we performed quantitative fluorescence intensity analyses of deconvolved confocal images taken through single myelin sheaths. This analysis revealed that the changes in fluorescence intensity along the length of a myelin sheath did indeed occur in quantal steps, suggesting differences in the number of wraps of myelin membrane along the axon, rather than random fluctuations of fluorescence signal intensity (Figure S1). When measuring how fluorescence intensities moved over time, we observed that along the axon not only the edges of the myelin segments extended laterally but also did the step-like fluorescence patterns (Figures 1A, 1B, and S1). Such dynamic changes in fluorescence intensity were only observed during the first few hours following initiation of myelin sheath formation. At later stages of myelin sheath maturation with a sheath length of more than 10 µm (observed using Tg(mbp:EGFP-CAAX)), the fluorescence intensity profiles along the length of individual sheaths were more complex, presumably because of an increased number of wraps and the exclusion of membrane-tethered GFP from the compact myelin sheaths (Figures 1D and 1E). Quantification of 25 mature and longer myelin sheaths showed that fluorescence intensity was on average uniform along the length of such mature sheaths (Figure 1F).

Thus, myelin growth appears to start nonuniformly at the center of nascent myelin sheaths and continues by lateral extension of developing layers toward the future nodes, which is consistent with a previous model of myelin growth (Sobottka et al., 2011).

### Three-Dimensional Reconstruction of Growing Myelin Sheaths

Because time-lapse imaging with light microscopy is not able to resolve single myelin layers, we used electron microscopy to investigate their spatial organization during myelination in mice. We first analyzed mouse optic nerves at postnatal day (P) 10, when myelin could be captured at all stages of growth with a minimal amount of mature sheaths. For sample preparation we used high-pressure freezing (HPF) followed by freeze substitution on fresh optic nerve to avoid artifacts typically introduced by chemical fixation.

To understand the spatial organization of a growing myelin sheath, we performed serial block-face imaging by focusedion beam milling in the SEM (FIB-SEM) to obtain high-resolution three-dimensional reconstructions of the optic nerve at P10 (Figures 2A-2C). The imaged volumes comprised 16 × 14 × 40–55  $\mu$ m (X, Y, Z) with a lateral resolution of 4–6 nm and a Z-resolution of 50 nm.

Within these volumes, we identified and reconstructed the cytoplasm-rich areas of the developing myelin sheath of 45 individual axons from three different animals to analyze the pattern of myelin growth. A representative three-dimensional reconstruction of such myelinated fibers is shown in Figure 2, and Movie S3 illustrates the typical features of a growing myelin sheath. We found that the thickest part of myelin is found where the process of the oligodendrocyte is connected to the myelin sheath. In addition, we observed that the outermost membrane layer spans the entire myelin segment length, with only minimal winding around the developing sheath (depicted in purple in Figures 2A, 2B, and 2F). In contrast, the three-dimensional reconstructions of the inner tongue clearly illustrate that it winds much more frequently around the axon as compared to the outer tongue. When following a myelinated segment from the thinnest part at the lateral edge (two wraps: Figure 2D) toward the thickest part at the center (11 wraps: Figure 2E), we observed that the average lateral distance between the windings of two successive layers was around 6 µm. A mirror pattern could be observed toward the other end of the sheath (Figures 2C and 2F). We could confirm these morphological features using ultrathin longitudinal sections of P10 optic nerve prepared for transmission electron microscopy (TEM), where individual membrane layers of myelin were discernible over an average distance of 20 µm (60 myelination events from three different animals) (Figure 2G). On longitudinal sections of the myelin sheath, the lateral endings of each individual layers appear as a cytoplasmic-rich edge. The distance between these lateral cytoplasmic-rich edges of the successive



### Figure 1. Live Imaging of Myelin Formation in Zebrafish

(A) Visualization of the growth of a myelin segment in live Tg(nkx2.2a:meGFP) zebrafish for 500 min at 3 days postfertilization (dpf). The steps in fluorescence intensity are pointed with arrowheads.

(B) Graphic representation of the relative fluorescence along the myelin segment over time shown in (A).

(C) Graph showing relative fluorescence intensity measurements for 32 nkx2.2a:mEGFP-expressing nascent myelin sheaths along their length. This indicates higher average intensity at the center of sheaths relative to their ends. Bars show mean ± SD.

(D) Graph showing length of 32 myelin sheaths imaged using the nkx2.2a:mEGFP compared to 25 segments visualized in the mbp:EGFP-CAAX zebrafish line.
(E) Lateral view of a mbp:EGFP-CAAX-expressing mature myelin sheath at 4 dpf shows fluorescence intensity pattern along the maturing myelin segment. Arrowheads point to areas of higher fluorescence intensity.

(F) Graph showing relative fluorescence intensity measurements for 25 mbp:EGFP-CAAX-expressing mature myelin sheaths along their length. Scale bars, 1 µm. See also Figure S1 and Movie S1.

layers along the length of the myelin sheath was  $\sim$ 5.2 µm in average (Figures 2G and S2). When we quantified the number of myelin layers relative to their distance from the oligodendrocyte process, we found that the number was highest at the position at which the oligodendrocyte process feeds into the myelin sheath and became lower with increasing distance at both edges of the growing myelin sheath (Figures 2C and 2G; Movie S4).

Next, we compared these early myelination profiles with later stages of development. When myelin structure was investigated on longitudinal sections at P23 and P60 (60 myelination profiles from three different animals), the distance of the lateral cytoplasmic-rich edges of the successive layers along the length of the myelin sheath increased from around 5.2  $\mu$ m at P10, to  $\sim$ 34.2  $\mu$ m at P23 and to  $\sim$ 137.3  $\mu$ m at P60 (Figure S2). Correspondingly, when the formation of the paranodes during the maturation of the myelin sheath was assessed, the distance between successive cytoplasmic-rich edges of the layers was

continuously declining with time until the paranodal loops were fully established at P60 (from  $\sim$ 2.5 µm at P10 to  $\sim$ 0.3 µm at P23 and  $\sim$ 0.2 µm at P60) (Figure S2).

Based on the shape of the myelin layers and their development over time our model suggests that a growing myelin sheath winds around the axon by an advancing inner tongue (underneath the previously deposited membrane) in the center of the myelin segment. Concurrent to this radial growth of the sheath, the layers extend laterally toward the node of Ranvier. This corresponds to our observations in live animals that membrane fluorescence intensity is at first higher at the center of a myelin sheath than at the lateral edges and evens with time.

### Polarized Transport of the Vesicular Stomatitis Virus G Protein to the Inner Tongue

EM reconstructions provide static images with high resolution, and time-lapse imaging using light microscopy is not able to



### Figure 2. Three-Dimensional Reconstruction of Myelin Structure in High-Pressure Frozen Optic Nerves

(A–C) High-resolution 3D reconstruction of a forming myelin segment showing the structure of the inner and outer layers (green, axolemma; orange, inner tongue; cyan, compacted myelin; purple, outer tongue). (B) The outer tongue is connected to the oligodendrocyte cell body (arrowhead), and the position at which the outer tongue has made one turn is marked by an arrow. (C) The inner tongue coils around the axon from a two-wraps thick myelin sheath (red line; cross-section shown in D) up to 11 wraps (white line; cross-section shown in E). Every turn is indicated by arrows pointing down and by arrows pointing up where the inner tongue is unwinding toward the other end of the sheath. Scale bar, 10 μm.

(D) Cross-sectional image of the area indicated with a red line (myelin with two wraps).

visualize adjacent myelin layers. We therefore sought out other methods to resolve at subcellular resolution, where newly synthesized membrane components are incorporated into the growing myelin sheath. To visualize the site at which proteins are integrated into the myelin sheath, a large amount of a membrane-associated reporter protein needs to be generated in a short time to allow for detection before the protein diffuses away into the growing sheath. Therefore, we used the vesicular stomatitis virus (VSV) to express glycoprotein G (VSV-G) as a tool, a well-established reporter of membrane protein trafficking within the biosynthetic pathway (Farguhar, 1985). To directly analyze the trafficking, first cultured oligodendrocytes were infected with VSV and analyzed at 2, 3, and 4 hr postinfection. At 2 hr postinfection VSV-G accumulated within the cell body and the cytoplasmic channels of the sheet in the Golgi-complex and in vesicular structures, but did not yet reach the membrane surface (Figure S3). Starting at 3 hr postinfection, VSV-G was detectable at the cell surface and, interestingly, showed enrichment at the F-actin-rich rim of the cell (Figure S3). The enrichment at the rim was not due to a larger membrane area, e.g., membrane folds, because the comparison to cells at 4 hr postinfection revealed that this enrichment diminished over time (Figure S3). Therefore, we conclude that insertion of VSV-G did not occur randomly in the cell but occurred selectively in the most distal region, from which the proteins can diffuse backward into the sheets.

We next investigated whether we could also use this system in vivo by localizing VSV-G or virus budding sites in myelinating oligodendrocytes. High-titer stocks of VSV were injected into the corpus callosum of P21 and P60 mice that were sacrificed 3 and 6 hr later. Immunoelectron microscopy at 3 and 6 hr postinfection showed that large amounts of VSV-G preferentially accumulated in the inner tongue membrane of the myelin sheath and much less in the outer tongue (Figures 3A, 3E, and S3). The antibody used was of high specificity, as no background staining could be observed in control-injected animals (Figure 3D). At 6 hr postinfection, this preferential distribution of the VSV-G at the inner tongue had increased even further (Figure S3). In addition, after 6 hr of infection, we observed an accumulation of viral particles at the inner tongue of the growing myelin sheath (Figure S3). In contrast, when VSV was injected into the corpus callosum of mice at P60 when myelin biogenesis is terminated, most of the labeling was found in the outer tongue, and there was hardly any VSV-G labeling at the inner tongue at 6 hr postinfection (Figures 3B and 3C).

Our results suggest that membrane trafficking in myelinating oligodendrocytes is polarized as new membrane protein is targeted and selectively inserted at the inner tongue of the growing myelin sheath.

### Cytoplasmic Channels Provide Direct Connections to the Leading Edge

A myelin growth zone at the inner tongue would require transport of newly synthesized membrane components from the soma through the oligodendroglial process. In principle, vesicles could traffic through cytoplasmic channels from the outside to the inside of the sheath along the helical coil that we have defined. In the peripheral nervous system (PNS) an additional connection of the outer and inner tongues of myelin is provided by the Schmidt-Lanterman incisures, but these structures have rarely been observed in CNS myelin (Blakemore, 1969; Hildebrand et al., 1993). However, in this study we used high-pressure freezing and freeze substitution to better preserve the native myelin structure for electron microscopy in order to analyze the prevalence of cytoplasmic channels. Surprisingly, using this method, we observed at the peak of myelination (P10) a large number of myelin sheaths with cytoplasmic channels, providing a short connection from the outer and inner tongue of myelin (Figures 4A-4C). There was a gradual reduction of these cytoplasmic channels with the further development of the myelin sheaths (Figure 4C). At P60 they had almost completely disappeared (Figure 4C). At P10 vesicular structures were frequently observed within the cytoplasmic channels, as well as in the inner and outer tongue, covering ~12% of the cytoplasmic area (Figure S4). Later in development, concomitant with the disappearance of the cytoplasmic channels, there was a dramatic decrease in the cytoplasmic area covered by vesicles (Figure S4). The three-dimensional reconstruction of a representative myelination profile at P10 shows a cytoplasmic channel system (Figure 4D, in red) that runs through a compact myelin sheath and ends at two points in the inner tongue (Figures 4E and 4F).

Furthermore, the three-dimensional reconstructions based on the serial block face imaging data set show that some cytoplasmic channels appear to run into and end within myelin outfoldings (Figures 4E and 4G). Myelin outfoldings (Figures 4G and 4H), also termed "redundant myelin," have previously been considered pathological features associated with different myelin diseases, including dysmyelinating neuropathies (Pereira et al., 2012). Here, we find that 6.5% of the myelinated axons had outfoldings in cross-section analysis of P10 optic nerve (Figure 4I). Based on the three-dimensional reconstruction with serial block face imaging, we determined that outfoldings have a width of around 10 µm; because a myelin sheath has an average length of around 100  $\mu$ m, we then estimated that at least half of the myelinated segments should have outfoldings at P10 in the optic nerve. Interestingly, myelin outfoldings almost completely disappeared at P60 (Figure 4I). The disappearance of the myelin outfodings correlated with the formation of the radial component by Claudin-11, a tight-junction component that fixes the adjacent layers of myelin membrane (Figure S4).

<sup>(</sup>E) Cross-sectional image of the area indicated with a white line (myelin with 11 wraps).

<sup>(</sup>F) Average distance between two lateral cytoplasmic-rich edges of successive myelin layers of the inner and outer tongue. Bars show mean ± SD (n = 3 animals with 45 different axons; \*\*\*p < 0,001, t test).

<sup>(</sup>G) Representative myelin sheath of P10 mouse optic nerve obtained by serial sectioning of high-pressure frozen tissue and imaged by TEM. An arrow is pointing to the lateral edge of successive myelin layers, each in a different color. (a–d) In the enlarged areas the individual myelin layers are marked by different colors and followed along the segment; the position of the zoomed areas is shown as a box. Scale bar, 5 μm.



P21 corpus callosum (control)





P21 corpus callosum 6 hours after VSV injection: longitudinal view

Figure 3. Tracking Membrane Trafficking Using the Vesicular Stomatitis Virus G Protein In Vivo

Ε

(A and B) High-titer stocks of VSV were injected into the corpus callosum of P21 and P60 mice. The subcellular localization of VSV-G in the myelin sheath was determined by immunoelectron microscopy 6 hr after infection at P21 (A) and P60 (B) mice (orange, inner tongue; purple, outer tongue). (C) Quantification of the VSV-G labeling distribution within the different domains of the myelin sheath at P21 and P60. Bars show mean ± SD (n = 3; 100 axon per animal; \*\*\*p < 0.001, t test).

(D) No background staining of the VSV-G antibody after control injections without virus into the corpus callosum at P21.

(E) Localization of VSV-G at the inner layers in longitudinal section of P21 corpus callosum 6 hr after VSV injection. Scale bars, 200 nm; gold size, 10 nm. See also Figure S3.

Together, these observations support the idea that the cytoplasmic channels are involved in facilitating the efficient delivery of membrane material to the active growth zone of the inner layers, which in some cases leads to transient focal hypermyelination as seen by the generation of myelin outfoldings (Figure S4).

### Regulation of Myelin Growth by PI(3,4,5)P3 In Vivo

Taken together, our data suggest that newly synthesized myelin membrane is directed through a system of cytoplasmic channels toward the inner tongue before fusing with the growing myelin sheath. If this is true, we should be able to affect the size and abundance of cytoplasmic channels by stimulating myelin growth. Indeed, the growth of the myelin sheath in vivo was associated with a larger inner but not a larger outer tongue when compared to later stages of development (Figures 5A and 5B). A key pathway that regulates myelin growth is the PI3K/AKT/mTOR signaling pathway (Flores et al., 2008; Tyler et al., 2009; Goebbels et al., 2010).

Thus, we analyzed whether PI(3,4,5)P3 elevation in vivo would influence the frequency of cytoplasmic channels and the size

of the inner tongue and, in so doing, promote myelin growth. Indeed, when myelin was analyzed in the optic nerves after conditional inactivation of Pten, which leads to PI(3,4,5)P3 elevation at P23 and P60, a much larger inner tongue and more cytoplasmic channels were observed as compared to the control, whereas the size of the outer tongue was unaffected (Figures 5C, 5D, and 5H-5J). In addition, when we investigated myelin structure on longitudinal sections at P23 and P60, we found that in contrast to wild-type animals, where the cytoplasmic-rich edges had assembled into paranodes, a large number of cytoplasmic-rich edges were still found along the length of the myelin sheath in Pten-deficient mice (Figures 5E and 5F). Furthermore, the area covered by vesicular profiles within the uncompacted areas of the myelin sheath was much higher in the Pten-deficient mice as compared to wild-type at P23 and P60 (Figure 5G). Thus, all of the key morphological features observed in the developing myelin sheaths of wild-type animals that are lost in mature adult sheaths reappear in the adult hypermyelinating Pten mutants, confirming a role of these features in myelin growth.

We also asked whether reinitiation of myelin growth involves the reopening of these channels after completion of developmental myelination. To test this, we analyzed *Pten*<sup>*flox/flox\*Plp1-CreERT2* mice 4 weeks after tamoxifeninduced conditional inactivation of *Pten* in oligodendrocytes at P100. We found in the optic nerve that a large number of channels reappeared (Figures 5K and 5L), at a time when the number of myelin wraps was unaltered (Figure 5M). To determine whether the opening of the channels was followed by an increase in myelin wrapping, we analyzed *Pten*<sup>*flox/flox\*Plp1-CreERT2* mice 12 weeks after tamoxifen injection. Indeed, we did observe that, in addition to an increase of cytoplasmic channels, myelin was significantly thicker, suggesting that reopening of the cytoplasmic channels precedes renewed myelin growth (Figures 5N and 5O). Thus, the cytoplasmic channels appear to be dynamic and have the potential to reopen after the developmental program is terminated.</sup></sup>

## Regulation of Membrane Trafficking by PI(3,4,5)P3 to the Leading Edge In Vitro

We next asked whether PI(3,4,5)P3 regulates membrane trafficking in oligodendrocytes. To this end, we turned to cell culture experiments and first confirmed that the size of the cellular rim of cultured oligodendrocytes increased during cellular growth and was enriched in F-actin, PI(3,4,5)P3 and Phospho-Akt (Figure S5). In order to verify that the outer rim of an oligodendrocyte in culture corresponds to the inner tongue of myelin in vivo, we analyzed whether there was a similar enrichment of F-actin in the inner tongue of the myelin sheath in the mouse during the active phase of myelination. Indeed, F-actin colocalized in a patch-like structure with MAG, which resides at the adaxonal membrane, but to a lesser extent with MBP, a marker of compact myelin (Figure S5), as described in the PNS (Trapp et al., 1989). In contrast, at P60 when the active growth phase of myelin has passed, F-actin was undetectable (Figure S5). Next, we determined whether the size of the outer rim/leading edge is influenced by the PI(3,4,5)P3/Akt signaling pathway. By applying a panel of well-characterized inhibitors or activators, we found that inhibition of PI(3,4,5)P3/Akt signaling resulted in thinning of the leading edge, whereas the activation using a PTEN inhibitor had the opposite effect (Figure S5). We then asked whether PI(3,4,5)P3 directly guides membrane trafficking in oligodendrocytes and applied a method to artificially polarize cell growth. Electrical signals are known to regulate cell migration by the activation of PTEN-dependent signaling and polarized elevation of PI(3,4,5)P3 (Zhao et al., 2006). When oligodendrocytes were cultured in a physiological electric field (1 V/cm for 2.5 hr), we observed that Phospho-Akt distribution was found polarized toward the cathode in 70% of the cells (Figure S5). When cells were infected with VSV and membrane trafficking was followed 3 hr postinfection, the targeting of VSV-G occurred preferentially toward the cathode (Figure S5). Thus, our results demonstrate that membrane trafficking of VSV-G is not random but polarized toward a growing edge in cultured oligodendrocytes.

## Regulation of Myelin Membrane Compaction by MBP and CNP

In addition to the membrane transport along the biosynthetic pathway, cytoplasmic components also need to be delivered

to myelin. MBP is the major cytoplasmic component of myelin and is known to mediate the compaction of the cytoplasmic leaflets of the myelin lamellae, a process that excludes most cytoplasm from the developing myelin sheath and which is necessary to provide myelin with the crucial physical stability for myelin membrane wrapping (Figure S6). As previously shown in multiple electron microscopy analysis of myelin (Hildebrand et al., 1993), compaction is fast and starts early in development after a few wraps. To analyze the process of myelin compaction in more detail, we investigated the extent of compaction at the age of 10, 14, and 21 days in the optic nerves of wild-type and heterozygote shiverer mice in which myelin compaction is expected to occur slower because of decreased MBP expression levels. In wild-type nerves studied at P10, 12% of myelinated axons were not yet fully "compacted" (Figures 6A and 6B). Our high-resolution EM analyses showed that compaction almost invariably appears first in the outermost layers of myelin (Figure S6), suggesting that the addition of new layers by the growth of the inner tongue occurs faster than the MBP-mediated compaction. Importantly, in optic nerves of heterozygote shiverer mice, more myelin sheaths with uncompacted wraps were detected. In addition, the number of compacted membrane layers in relation to the total number of myelin layers was lower compared to wild-type at P10 and P14 (Figure S6). At the age of P21, no significant differences were detectable between these aenotypes (Figure 6B).

MBP mRNA is transported to the growing tip of oligodendroglial processes (Ainger et al., 1993), and translation of MBP is induced at the innermost myelin layers that are in contact with the axon (Colman et al., 1982; Trapp et al., 1987; Wake et al., 2011; Laursen et al., 2011). However, when we analyzed the subcellular distribution of MBP protein in the myelin sheath by immunoelectron microscopy, the levels of MBP were 4-fold lower in the inner uncompacted wraps as compared to the outer compacted layers (Figures 6C and 6D). Thus, we hypothesized that there must be active mechanisms that prevent myelin compaction at the innermost regions of myelin. Because these noncompacted wraps were enriched in CNP (Figure S6), we analyzed their role in this process. Indeed, when the optic nerves of heterozygote and homozygotes CNP deficient mice were examined by high-pressure freezing EM at P10, a dramatic reduction of the uncompacted myelin wraps were observed. Whereas ~12% of myelin sheaths contained uncompacted wraps in wild-type mice, there was a reduction to  ${\sim}4\%$  in CNP +/- and to less than 1% in CNP -/- mice (Figure 6E). Thus, our results show that the equilibrium of CNP and MBP levels appears to regulate the speed of compaction in early development. Hence, we conclude that the rapid extension of the innermost tongue during myelin growth, together with the accumulation of CNP between these newly synthesized layers, prevent the premature compaction of myelin by MBP at the leading edge.

### DISCUSSION

The aim of this study was to answer a fundamental question of axon-glia interactions in the vertebrate nervous system: how does an oligodendrocyte "wrap" its plasma membrane around



Figure 4. Cytoplasmic Channels and Myelin Outfoldings Appear Transiently in Developing Myelin Sheaths

(A and B) Longitudinal and cross-sections of P10 optic nerve myelinated axons showing cytoplasmic channels within the compact myelin indicated by arrow heads. Scale bar, 200 nm.

(C) Percent of myelin sheaths with cytoplasmic channels at P10, P14, P21, and P60. Bars show mean  $\pm$  SD (n = 3; 200 axons per animal; \*p < 0.05, \*\*p < 0.01, t test).

Cell

the axon? In this study we have combined data from very different experimental approaches that have led us to a model of myelin membrane growth in the central nervous system, which is summarized in Figures 7, S7, and Movie S5: we suggest that myelin grows by two coordinated motions: the wrapping of the leading edge at the inner tongue in a triangle-like shape around the axon underneath the previously deposited membrane, together with the lateral extension of myelin membrane layers toward the nodal regions. In this model, the lateral cytoplasmic domains of each myelin layer are always in contact with the axonal surface and move in a continuous helical manner toward the future node, where they align and form the paranodal loops. An elaborated system of cytoplasmic channels within compacted myelin provides shortcuts for the transport of membrane to the growth zone at the inner tongue in the developing myelin sheath. Most channels appear transiently in development at the most intense phase of membrane growth and are regulated by PI(3,4,5)P3 levels.

We used electron microscopy of high-pressure frozen samples and three-dimensional reconstruction based on serial block face to obtain snapshots of myelin ultrastructure during development of the optic nerve. We find that the innermost layer of myelin has the shortest lateral extension, whereas the layers toward the top of the myelin sheath build up in size with each consecutive wrap. Previous studies using light microscopy have observed a coil of an average periodicity of 5.7 to 7 µm along the internodal dimension (Pedraza et al., 2009; Sobottka et al., 2011; loannidou et al., 2012). This has led to the suggestion that myelin extends as a coil, twisting across the axon in a corkscrew motion (Pedraza et al., 2009), or that myelin thickening is achieved by forming new layers on to top of the inner ones in a "croissant-like" manner (Sobottka et al., 2011). One limitation of these studies is that they are almost entirely based on light microscopy. However, in our three-dimensional EM reconstructions we neither detect a serpent-like process coiling around the axon nor a "croissant-like" structure. A possible explanation for the conclusions of the previous studies could be the time point at which myelination was analyzed. In the very initial stage, the myelin layers may indeed show overlapping profiles at the surface of the forming sheath. However, we find that during myelin growth, the outermost myelin layer is continuous along the entire length of the sheath. The pattern of the coiling helix arises from the organization of the inner layers of myelin within the myelin sheath along the axon. In addition, and as has been noted previously (Webster, 1971; Knobler et al., 1976), the nonuniform thickness of the growing myelin internode is also incompatible with a "carpet crawler"-type of model in which myelin grows uniformly around the axon. A major limitation of all of the previously proposed models is that the leading edge was not defined. By using a virus as a reporter, we were able to follow membrane trafficking within the forming myelin sheath and define the inner tongue as the growth zone.

In addition, to membrane transport via the biosynthetic secretory pathway, cytoplasmic components need to be delivered to myelin. MBP is the major cytoplasmic component, and its mRNA is transported in cytoplasmic granules to myelin, where the local translation of MBP is initiated and the compaction of the cytoplasmic surfaces triggered (Colman et al., 1982). We observed that the targeting and the deposition of myelin membrane components are spatially separated in the growing myelin sheath. Whereas vesicular transport is directed toward the inner tongue of the myelin sheath, MBP-mediated compaction occurs first in the outer layers and moves inward with time. We propose that even if MBP is synthesized in the front, compaction is not fast enough to keep up with the rapid extension of the innermost tongue during myelin growth. In addition, we identified CNP, a protein previously implicated in maintaining axonal integrity (Lappe-Siefke et al., 2003), as a factor that delays myelin compaction. In the absence of CNP, myelin compaction appeared to proceed faster and extend to the inmost layers of the myelin sheath. These data are consistent with previous work in a transgenic mouse line, showing that overexpression of CNP results in areas that lack myelin compaction (Gravel et al., 1996). Taken together, we hypothesize that MBP is synthesized in the distal (close to the axon) areas of myelin, followed by its diffusion backward to the outermost area at which compaction is initiated. Once MBP is bound to the two adjacent cytoplasmic surfaces, it polymerizes into fibrous network, thereby greatly reducing its mobility and establishing a nucleation point for myelin membrane zippering (Aggarwal et al., 2013). This fusion of the adjacent cytoplasmic surfaces of the myelin bilayer is necessary to provide myelin with the crucial physical stability for myelin membrane wrapping. We hypothesize that myelin wrapping is maintained by tightly regulating growth and compaction together to ensure that a stable (Toyama et al., 2013) and tightly condensed membrane is driven multiple times around the axon.

Another important question is how membrane trafficking is polarized toward the leading edge. PI3-kinase activation and its lipid product PI(3,4,5)P3 are known to play instructive roles in cell polarization by recruiting and activating the small RhoGTPases at the leading edge (Ridley, 2011). Axonal signal may also regulate myelin growth. Electrical activity, for example, influences myelination in the CNS, possibly by effectors, such as neurotransmitters, ATP, ions, or growth factors (Barres and Raff, 1993; Demerens et al., 1996; Fields and Stevens-Graham, 2002).

t test). See also Figure S4.

<sup>(</sup>D) Three-dimensional organization of the cytoplasmic channels (red, cytopalsmic channels; orange, inner tongue; green, axolemma) along the myelin sheath (thick arrow signals the location the oligodendrocyte process). Scale bar, 10  $\mu$ m.

<sup>(</sup>E and F) Model and electron microscopy views of the area at which the cytoplasmic channels (red) reach the inner tongue (arrowhead: position of the EM views). Scale bar, 500 nm.

<sup>(</sup>G) Spatial organization of cytoplasmic channels (red) running inside the outfolding and reaching the inner tongue at the lateral end of the outfolding (red arrow head).

<sup>(</sup>H) Cross-section morphology of myelin outfolding (arrowheads) in P10 high-pressure frozen optic nerve (star labels the axon with a normal myelin sheath). (I) Amount of myelin outfoldings in myelinated fibers in optic nerves between P10 and P60. Bars show mean  $\pm$  SD (n = 3; 200–800 axons per animal; \*p < 0.05,

### Cell



(legend on next page)

We found that an electrical field was sufficient to polarize the leading edge and membrane trafficking toward the cathode. Such electrical fields might be generated by the local accumulation of  $K^+$  when released from spiking axons.

Another way to guide membrane trafficking to the leading edge is provided by the cytoplasmic channels with the myelin sheath. The detection of numerous cytoplasmic channels in developing myelin was unexpected, as Schmidt-Lantermann incisures-like structures were not thought to be abundant in CNS myelin (Bunge, 1968). An important technical advancement that made the visualization of cytoplasmic channels possible was highpressure freezing of fresh tissue for electron microscopy, which results in preservation of tissue architecture close to native state. Injection of fluorescent dyes into freshly isolated spinal cord slices has also revealed an extensive network of interconnected cytoplasmic pockets in myelin (Velumian et al., 2011).

Recent studies suggest that activity dependent myelination should not only be considered in development (Wake et al., 2011) but also be considered in nervous system plasticity in postnatal development and adulthood (Liu et al., 2012; Makinodan et al., 2012; Young et al., 2013). Notably, social isolation reportedly impairs myelination in the prefrontal cortex in young mice, whereas social reintegration is sufficient to normalize myelin formation (Liu et al., 2012). Consistent with the concept of adult myelin plasticity, myelin growth can be experimentally reactivated even if the developmental program has been terminated (Goebbels et al., 2010). Here, we show that the cytoplasmic channels reopen when PI(3,4,5)P3 levels are increased in the adult mice, which was followed by an increase in myelin thickness. Thus, this elaborated network of interconnected tunnels is dynamic and provides a means to transport molecules through the internode even in fully developed myelin sheaths. One interesting question is how the formation of the cytoplasmic channels is regulated. An attractive hypothesis is that PI(3,4,5)P3 induces local actin polymerization at the cytoplasmic leaflet of the myelin bilayer, which covers and protects myelin from being compacted by MBP. It is possible that there is a tug-of-war-type regulation of the cytoplasmic channels by PI(3,4,5)P3 effectors, such as the actin cytoskeleton, possibly in association with CNP, holding MBP from membrane compaction.

We show that some cytoplasmic channels end within the myelin outfoldings, and we therefore propose that myelin outfoldings form by the preferential delivery of membrane through cytoplasmic channels into the inner tongue, resulting in focal accumulation of excess membrane that extends outward. These myelin outfoldings have previously been considered pathological features associated with different myelin diseases, including dysmyelinating neuropathies. Here, however, we clearly show that myelin outfoldings are part of the normal development of myelin in the optic nerve. When cytoplasmic channels close, these outfoldings may slowly spread out because of the fluidity of the membrane. Thus, outfoldings may serve as membrane reservoirs to allow the completion of the lateral growth of the layers and secure unperturbed radial growth of the myelinated axon itself, which would explain their gradual disappearance later in life.

Taken together, we have used an in-depth, integrated approach of advanced imaging at the light and electron microscopic level to dissect the morphogenesis of myelin membrane wrapping. Combined with the discovery of the previously unknown growth zone of myelin and of transient cytoplasmic channels in myelin that allow membrane trafficking toward the growing tip, our observations lead to a model of myelin wrapping. This model not only fills a gap in our understanding of myelin biogenesis in brain development but also sheds light on myelin pathology in various neurological diseases in which myelination is perturbed.

#### **EXPERIMENTAL PROCEDURES**

Full details are provided in the Extended Experimental Procedures.

#### **Fish Husbandry**

The transgenic zebrafish lines Tg(nkx2.2a:mEGFP) and Tg(mbp:EGFP-CAAX) were used in this study. The imaging is described in the Extended Experimental Procedures.

### **Mice and Primary Cell Culture**

The generation of PTEN cre/+ flox/floxPten and CNP mutant mice has been described previously (Goebbels et al., 2010; Lappe-Siefke et al., 2003). The *shiverer* mice lacking MBP were maintained on a C57Bl6 background. The

Figure 5. Regulation of Cytoplasmic Channels, Inner Tongue Size, and Myelin Sheath Growth by PI(3,4,5)P3 Levels

(A) Electron micrograph of high-pressure frozen wild-type optic nerve showing larger inner tongue at P14 (stars) as compared to P60. Scale bar, 200 nm. (B) Relative size of the inner and outer tongue area to the axonal area between P10 and P60.

(C) Electron micrograph of P60 conditional Pten-deficient mice (Pten KO; Pten<sup>flox/flox</sup> \*Cnp1-Cre/+) optic nerve showing enlarged inner tongues (stars).

(D) Relative size of the inner and outer tongue area to the axonal area at P23 and P60 for the Pten mutant (wild-type P60 data from B for comparison).

(E) Longitudinal view of P60 *Pten* mutant optic nerve shows the short lateral distances between successive inner layers (white arrow heads) and the large cytoplasmic channels within the compact myelin (black arrow heads).

(F) Average distance between two lateral cytoplasmic-rich edges of successive myelin layers at P23 and P60 for the *Pten* mutant compared to wild-type. (G) Quantification of the area covered by vesicular structures in *Pten*-deficient animals as compared to wild-type at P23 and P60 (data used for wild-type are from Figure S4).

(H) Electron micrograph of myelin from conditional *Pten*-deficient mice (*Pten* KO; *Pten*<sup>flox/flox</sup> \**Cnp*1-*Cre*/+) optic nerve compared to the control (Control; *Pten*<sup>flox/wt</sup> \**Cnp*1-*Cre*/+) at P60 (arrowheads pointing at colored cytoplasmic channels).

(I and J) Amount of myelin sheaths with cytoplasmic channels in cross-sections and measurement of the myelin thickness in control and *Pten* mutant optic nerves. (K) Electron micrograph of optic nerve myelin after tamoxifen-induced conditional inactivation for 4 weeks of *Pten<sup>flox/flox</sup>\*Plp1-CreERT2* mice in the adult (P60) compared to the control (arrowhead pointing at colored cytoplasmic channels). Scale bar, 500 nm.

(L–O) Amount of myelin sheaths with cytoplasmic channels in cross-sections and measurement of the myelin thickness in control and after tamoxifen-induced conditional inactivation of *Pten* for 4 weeks (L and M) and for 12 weeks (N and O). Bars shown in all graphs are the mean  $\pm$  SD (n = 3; 100–200 axons per animal; 20 axons per animals for F; \*p < 0.05, \*\*p < 0.01, \*\*\*p < 0.001, t test). See also Figure S5.



Shiv CNP CNP

+/-+/-

-/-

WТ

primary oligodendrocyte cell culture was prepared as described previously (Aggarwal et al., 2013).

### Electron Microscopy

0.00

Mice were killed by cervical dislocation, and freshly extracted optic nerves were cryofixed using a high-pressure freezer HPM100 (Leica) and further processed by freeze substitution and EPON-embedding following the "tannic acid-OsO<sub>4</sub> protocol" described in Möbius et al. (2010). Cross- and longitudinal ultrathin sections were obtained with an Ultracut S ultramicrotome (Leica) and contrasted as described previously (Möbius et al., 2010). Sections were imaged using a LEO 912 Omega electron microscope (Zeiss) equipped with an on-axis 2k CCD camera (TRS). Three to five animals were used for each analysis. On cross-sections, five to ten randomly selected areas of 150  $\mu\text{m}^2$ were imaged per animal in which 100 to 300 myelinated axon profiles with four or more myelin wraps were counted. For the assessment of the number

### Figure 6. Myelin Compaction from the Outer to the Inner Layers Is Regulated by MBP and CNP Levels

(A) Cross-section of P10 wild-type myelin sheath with outer layers compacted (black arrows) and inner layers noncompacted (white arrows).

(B) Amount of myelinated axons with noncompacted lavers in wild-type and shiverer heterozygote (Shiv  $\pm$ ) optic nerves between P10 and P21. (C) Immunoelectron micrograph for MBP of a partially compacted myelin sheath of P10 wild-type optic nerve (black arrows pointing at compacted lavers and white arrows to the noncompacted lavers). Scale bars: 200 nm: gold size. 10 nm.

(D) Quantification of MBP labeling per  $\mu m$  of membrane in the compacted and noncompacted layers of P10 wild-type myelin sheaths.

(E) Amount of myelinated axons with noncompacted layers in wild-type, Shiv ±, CNP ±, and CNP -/- at P10. Bars show mean ± SD (n = 3–5; 120–200 axons per animal; \*\*p < 0.01, \*\*\*p < 0.001, t test).

See also Figure S6.

of wraps for shiverer -/- and control all myelinated optic nerves were analyzed. For morphological analysis on longitudinal sections, at least 20 myelinated axonal segments of 10-30 µm were assessed per animal.

Preparation of optic nerves for immunoelectron microscopy and immunolabeling of Tokuyasu sections of 50 nm were performed as described previously (Werner et al., 2007). In brief, optic nerves were freshly removed and immersion fixed in 4% formaladehyde, 0.25% glutaraldehyde (EM grade, Science Services) in phosphate buffer (pH 7.3) as described in Möbius et al. (2010). After a few hours of fixation, the nerves were embedded in 10% gelatin (Twee Torens), infiltrated in 2.3 M sucrose overnight, mounted onto aluminum pins, and frozen in IN2. Tokuyasusections were prepared with a UC6 cryoultramicrotome (Leica) and a cryoimmuno diamond knife (Diatome). For spinal cord samples, mice at the age of 4 and 60 days were perfused transcardially with the same fixative described above. Then the spinal cord was carefully dissected and postfixed for 12 hr in 4% paraformaldehyde. The thoracic spinal cord portion was extracted from the bones and further pro-

cessed for immunoelectron microscopy. We collected 400 nm thick Tokuyasu sections of the spinal cord ventral column on a super frost microscopic slice and processed them for immunofluorescence staining.

Immunolabeling of ultrathin (50 nm) Tokuyasu sections was performed as described in Peters and Pierson (2008) using antibodies directed against (VSV-G, MBP, and CNP). Rabbit polyclonal antibodies were detected using protein A gold (10 nm, CMC). Monoclonal antibodies were also detected with protein A gold after incubation with a bridging antibody raised in rabbit. The quantification of the VSV-G labeling was performed in the corpus callosum on 100 axons wrapped by infected myelin sheath in three littermate animals for the two following conditions: 3 and 6 hr after VSV injection.

### Serial Block Face Imaging and 3D Modeling Software

Serial block face imaging data were obtained on a Helios Nanolab 650 dual beam scanning electron microscope equipped with a focused ion beam,



### Figure 7. Model of Myelin Biogenesis in the CNS

(A–D) Model of a developing myelin sheath in wrapped, unwrapped, and cross-section views. The unwrapped representation shows the geometry and the development of the sheath and the localization of the cytoplasmic channels, which connect the cell body and the growth zone at the inner tongue. The growth zone is colored in pink, and the compacted myelin is in dark violet. The wrapped representation shows the position of the layers when wrapped around the axon. The crosssections show the state of compaction during myelin growth.

See also Figure S7 and Movie S5.

and images were obtained with a 2.0 kV electron beam using the back scatter electron (BSE) mode (immersion lens). The block faces were imaged using the slice and view software with a dwell time of 3  $\mu$ s. Total volumes of 16  $\times$  13.8  $\times$  40–55  $\mu$ m<sup>3</sup> were recorded with a x/y pixel size of 4–6 nm during 4–7 days of acquisition in each sample. The alignment, segmentation, and three-dimensional (3D) reconstruction was performed using the Amira and the IMOD software (Kremer et al., 1996). Models derived from the experimental data were drawn in the Corel Draw 5 software suite.

#### **VSV** Injections

For intracerbral (i.c.) infection, mice were anesthetized using isoflurane inhalation anesthesia. i.c. inoculation (10  $\mu$ l) was performed using a 28G insulin syringe over a period of 3 min. Solution contains VSV in balanced salt solution and 0.3% of monostral blue. The latter was added for better visualization of the injection side for histological processing. Following i.c. injection, animals were euthanized either 3 or 6 hr later by intraperitoneal injection of pentobarbital and transcardially perfused with 4% PFA. Subsequently, brains were dissected, and small pieces of cortex with corpus callosum were postfixed with PFA for 4 hr and further processed for immunoelectron microscopy.

### SUPPLEMENTAL INFORMATION

Supplemental Information includes Extended Experimental Procedures, seven figures, and five movies and can be found with this article online at http://dx. doi.org/10.1016/j.cell.2013.11.044.

#### ACKNOWLEDGMENTS

We thank Torben Ruhwedel for technical assistance. We thank Ueli Suter for providing *Plp1-CreERT2* mice. This work was supported by a European Research Council (ERC) Starting Grant (to M.S.), grants from the German Research Foundation (SI 746/9-1; TRR43), E-Rare Research Program, the Tschira-Stiftung, a David Phillips Fellowship from the Biotechnology and Biological Sciences Research Council, a Research Prize from the Lister Institute, an International Reintegration Grant (to D.A.L.), and an EMBO Long-Term Fellowship (to T.C.). D.M. holds stipendiary professorships of the Swiss National Science Foundation (No. PP00P3\_128372). K.A.N. is supported by the German Research Foundation Research Center Molecular Physiology of the Brain (CMPB) and an ERC Advanced Investigator Grant (Axoglia).

Received: April 19, 2013 Revised: October 5, 2013 Accepted: November 7, 2013 Published: January 16, 2014

#### REFERENCES

Aggarwal, S., Snaidero, N., Pähler, G., Frey, S., Sánchez, P., Zweckstetter, M., Janshoff, A., Schneider, A., Weil, M.T., Schaap, I.A., et al. (2013). Myelin membrane assembly is driven by a phase transition of myelin basic proteins into a cohesive protein meshwork. PLoS Biol. *11*, e1001577.

Ainger, K., Avossa, D., Morgan, F., Hill, S.J., Barry, C., Barbarese, E., and Carson, J.H. (1993). Transport and localization of exogenous myelin basic protein mRNA microinjected into oligodendrocytes. J. Cell Biol. *123*, 431–441.

Barres, B.A., and Raff, M.C. (1993). Proliferation of oligodendrocyte precursor cells depends on electrical activity in axons. Nature 361, 258–260.

Ben Geren, B. (1954). The formation from the Schwann cell surface of myelin in the peripheral nerves of chick embryos. Exp. Cell Res. 7, 558–562.

Blakemore, W.F. (1969). Schmidt-Lantermann incisures in the central nervous system. J. Ultrastruct. Res. 29, 496–498.

Bunge, R.P. (1968). Glial cells and the central myelin sheath. Physiol. Rev. 48, 197–251.

Bunge, M.B., Bunge, R.P., and Ris, H. (1961). Ultrastructural study of remyelination in an experimental lesion in adult cat spinal cord. J. Biophys. Biochem. Cytol. *10*, 67–94.

Bunge, R.P., Bunge, M.B., and Bates, M. (1989). Movements of the Schwann cell nucleus implicate progression of the inner (axon-related) Schwann cell process during myelination. J. Cell Biol. *109*, 273–284.

Colman, D.R., Kreibich, G., Frey, A.B., and Sabatini, D.D. (1982). Synthesis and incorporation of myelin polypeptides into CNS myelin. J. Cell Biol. *95*, 598–608.

Czopka, T., Ffrench-Constant, C., and Lyons, D.A. (2013). Individual oligodendrocytes have only a few hours in which to generate new myelin sheaths in vivo. Dev. Cell *25*, 599–609.

Demerens, C., Stankoff, B., Logak, M., Anglade, P., Allinquant, B., Couraud, F., Zalc, B., and Lubetzki, C. (1996). Induction of myelination in the central nervous system by electrical activity. Proc. Natl. Acad. Sci. USA *93*, 9887–9892. Farquhar, M.G. (1985). Progress in unraveling pathways of Golgi traffic. Annu. Rev. Cell Biol. 1, 447–488.

Fields, R.D., and Stevens-Graham, B. (2002). New insights into neuron-glia communication. Science 298, 556–562.

Flores, A.I., Narayanan, S.P., Morse, E.N., Shick, H.E., Yin, X., Kidd, G., Avila, R.L., Kirschner, D.A., and Macklin, W.B. (2008). Constitutively active Akt induces enhanced myelination in the CNS. J. Neurosci. *28*, 7174–7183.

Fünfschilling, U., Supplie, L.M., Mahad, D., Boretius, S., Saab, A.S., Edgar, J., Brinkmann, B.G., Kassmann, C.M., Tzvetanova, I.D., Möbius, W., et al. (2012). Glycolytic oligodendrocytes maintain myelin and long-term axonal integrity. Nature *485*, 517–521.

Goebbels, S., Oltrogge, J.H., Kemper, R., Heilmann, I., Bormuth, I., Wolfer, S., Wichert, S.P., Möbius, W., Liu, X., Lappe-Siefke, C., et al. (2010). Elevated phosphatidylinositol 3,4,5-trisphosphate in glia triggers cell-autonomous membrane wrapping and myelination. J. Neurosci. *30*, 8953–8964.

Gravel, M., Peterson, J., Yong, V.W., Kottis, V., Trapp, B., and Braun, P.E. (1996). Overexpression of 2',3'-cyclic nucleotide 3'-phosphodiesterase in transgenic mice alters oligodendrocyte development and produces aberrant myelination. Mol. Cell. Neurosci. 7, 453–466.

Hildebrand, C., Remahl, S., Persson, H., and Bjartmar, C. (1993). Myelinated nerve fibres in the CNS. Prog. Neurobiol. *40*, 319–384.

Ioannidou, K., Anderson, K.I., Strachan, D., Edgar, J.M., and Barnett, S.C. (2012). Time-lapse imaging of the dynamics of CNS glial-axonal interactions in vitro and ex vivo. PLoS ONE 7, e30775.

Jessen, K.R., and Mirsky, R. (2005). The origin and development of glial cells in peripheral nerves. Nat. Rev. Neurosci. 6, 671–682.

Kirby, B.B., Takada, N., Latimer, A.J., Shin, J., Carney, T.J., Kelsh, R.N., and Appel, B. (2006). In vivo time-lapse imaging shows dynamic oligodendrocyte progenitor behavior during zebrafish development. Nat. Neurosci. *9*, 1506–1511.

Knobler, R.L., Stempak, J.G., and Laurencin, M. (1976). Nonuniformity of the oligodendroglial ensheathment of axons during myelination in the developing rat central nervous system. A serial section electron microscopical study. J. Ultrastruct. Res. *55*, 417–432.

Kremer, J.R., Mastronarde, D.N., and McIntosh, J.R. (1996). Computer visualization of three-dimensional image data using IMOD. J. Struct. Biol. *116*, 71–76.

Lappe-Siefke, C., Goebbels, S., Gravel, M., Nicksch, E., Lee, J., Braun, P.E., Griffiths, I.R., and Nave, K.A. (2003). Disruption of Cnp1 uncouples oligodendroglial functions in axonal support and myelination. Nat. Genet. *33*, 366–374.

Laursen, L.S., Chan, C.W., and Ffrench-Constant, C. (2011). Translation of myelin basic protein mRNA in oligodendrocytes is regulated by integrin activation and hnRNP-K. J. Cell Biol. *192*, 797–811.

Lee, Y., Morrison, B.M., Li, Y., Lengacher, S., Farah, M.H., Hoffman, P.N., Liu, Y., Tsingalia, A., Jin, L., Zhang, P.W., et al. (2012). Oligodendroglia metabolically support axons and contribute to neurodegeneration. Nature *487*, 443–448.

Liu, J., Dietz, K., DeLoyht, J.M., Pedre, X., Kelkar, D., Kaur, J., Vialou, V., Lobo, M.K., Dietz, D.M., Nestler, E.J., et al. (2012). Impaired adult myelination in the prefrontal cortex of socially isolated mice. Nat. Neurosci. *15*, 1621–1623.

Makinodan, M., Rosen, K.M., Ito, S., and Corfas, G. (2012). A critical period for social experience-dependent oligodendrocyte maturation and myelination. Science 337, 1357–1360.

Pedraza, L., Huang, J.K., and Colman, D. (2009). Disposition of axonal caspr with respect to glial cell membranes: Implications for the process of myelination. J. Neurosci. Res. *87*, 3480–3491.

Möbius, W., Cooper, B., Kaufmann, W.A., Imig, C., Ruhwedel, T., Snaidero, N., Saab, A.S., and Varoqueaux, F. (2010). Electron microscopy of the mouse central nervous system. Methods Cell Biol. *96*, 475–512.

Peters, P.J., and Pierson, J. (2008). Immunogold labeling of thawed cryosections. Methods Cell Biol. 88, 131–149.

Pereira, J.A., Lebrun-Julien, F., and Suter, U. (2012). Molecular mechanisms regulating myelination in the peripheral nervous system. Trends Neurosci. 35, 123–134.

Ridley, A.J. (2011). Life at the leading edge. Cell 145, 1012–1022.

Salzer, J.L., Brophy, P.J., and Peles, E. (2008). Molecular domains of myelinated axons in the peripheral nervous system. Glia 56, 1532–1540.

Sherman, D.L., and Brophy, P.J. (2005). Mechanisms of axon ensheathment and myelin growth. Nat. Rev. Neurosci. *6*, 683–690.

Sobottka, B., Ziegler, U., Kaech, A., Becher, B., and Goebels, N. (2011). CNS live imaging reveals a new mechanism of myelination: the liquid croissant model. Glia 59, 1841–1849.

Toyama, B.H., Savas, J.N., Park, S.K., Harris, M.S., Ingolia, N.T., Yates, J.R., 3rd, and Hetzer, M.W. (2013). Identification of long-lived proteins reveals exceptional stability of essential cellular structures. Cell *154*, 971–982.

Trapp, B.D., Moench, T., Pulley, M., Barbosa, E., Tennekoon, G., and Griffin, J. (1987). Spatial segregation of mRNA encoding myelin-specific proteins. Proc. Natl. Acad. Sci. USA *84*, 7773–7777.

Trapp, B.D., Andrews, S.B., Wong, A., O'Connell, M., and Griffin, J.W. (1989). Co-localization of the myelin-associated glycoprotein and the microfilament components, F-actin and spectrin, in Schwann cells of myelinated nerve fibres. J. Neurocytol. *18*, 47–60.

Tyler, W.A., Gangoli, N., Gokina, P., Kim, H.A., Covey, M., Levison, S.W., and Wood, T.L. (2009). Activation of the mammalian target of rapamycin (mTOR) is essential for oligodendrocyte differentiation. J. Neurosci. *29*, 6367–6378.

Velumian, A.A., Samoilova, M., and Fehlings, M.G. (2011). Visualization of cytoplasmic diffusion within living myelin sheaths of CNS white matter axons using microinjection of the fluorescent dye Lucifer Yellow. Neuroimage *56*, 27–34.

Wake, H., Lee, P.R., and Fields, R.D. (2011). Control of local protein synthesis and initial events in myelination by action potentials. Science 333, 1647–1651.

Webster, H.D. (1971). The geometry of peripheral myelin sheaths during their formation and growth in rat sciatic nerves. J. Cell Biol. *48*, 348–367.

Werner, H.B., Kuhlmann, K., Shen, S., Uecker, M., Schardt, A., Dimova, K., Orfaniotou, F., Dhaunchak, A., Brinkmann, B.G., Möbius, W., et al. (2007). Proteolipid protein is required for transport of sirtuin 2 into CNS myelin. J. Neurosci. 27, 7717–7730.

Young, K.M., Psachoulia, K., Tripathi, R.B., Dunn, S.J., Cossell, L., Attwell, D., Tohyama, K., and Richardson, W.D. (2013). Oligodendrocyte dynamics in the healthy adult CNS: evidence for myelin remodeling. Neuron *77*, 873–885.

Zhao, M., Song, B., Pu, J., Wada, T., Reid, B., Tai, G., Wang, F., Guo, A., Walczysko, P., Gu, Y., et al. (2006). Electrical signals control wound healing through phosphatidylinositol-3-OH kinase-gamma and PTEN. Nature 442, 457–460.